

MY ENTYVIO JOURNEY

AT A GLANCE

Ready to make the most of your ENTYVIO journey? Though no two paths are the same, here are the key steps you should know as you get started on ENTYVIO.

Got questions about where you are in your ENTYVIO journey? Give us a call at **1-844-ENTYVIO** (1-844-368-9846).



STARTING TREATMENT

STEP 1 GET PRESCRIBED ENTYVIO

[GO TO PAGE 2](#)

STEP 2 RECEIVE THE ENTYVIO PATIENT STARTER KIT

STEP 3 SIGN UP FOR *ENTYVIOCONNECT*

[GO TO PAGE 3](#)

STEP 4 STARTER IV DOSES

STEP 5 MAINTENANCE EVALUATION

The rest of your ENTYVIO journey will depend on whether you and your healthcare provider have decided to continue with intravenous (IV) infusions or self-administered injections under the skin (subcutaneous).

[GO TO PAGE 4](#)

IV INFUSIONS

STEP 6 FIRST IV MAINTENANCE DOSE

[GO TO PAGE 5](#)

STEP 7 ONGOING IV MAINTENANCE

SELF-ADMINISTERED INJECTIONS

STEP 6 RECEIVE SELF-INJECTION TRAINING KIT

[GO TO PAGE 6](#)

STEP 7 FIRST SELF-INJECTION MAINTENANCE DOSE

STEP 8 ONGOING SELF-INJECTION MAINTENANCE

What is ENTYVIO?

ENTYVIO is a prescription medicine used in adults for the treatment of:

- moderately to severely active ulcerative colitis (UC).
- moderately to severely active Crohn's disease (CD).

Safety Considerations

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 7](#).



1 GET PRESCRIBED ENTYVIO

Every ENTYVIO journey starts with a prescription from your healthcare provider. Make sure to use this time to ask any questions you have about receiving ENTYVIO.



What should I do now?

Schedule your first infusion. This could happen at a healthcare provider's office, an infusion center, or your home.

2 RECEIVE THE ENTYVIO PATIENT STARTER KIT

This kit has helpful information and resources for getting started on ENTYVIO. If you didn't receive your kit when you were prescribed ENTYVIO, talk to your healthcare provider.



What should I do now?

Dig into all the resources in your kit and online at www.entyvio.com/journey. The more you know, the more confident you'll be.

Safety Considerations

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 7](#).



3

SIGN UP FOR *EntyvioConnect*

EntyvioConnect offers resources to help you after you've been prescribed ENTYVIO, such as:

- Nurse Educators
- Financial and insurance help
- Ongoing support
- Self-Injection Training Kit



For more information and to sign up for *EntyvioConnect*, visit www.entyvio.com/connect

What should I do now?

Get in touch with your *EntyvioConnect* Nurse Educator. They can provide you with resources and information but cannot offer medical advice.

4

STARTER IV DOSES

Everyone who starts on ENTYVIO will have 2 or more starter doses by intravenous (IV) infusion.


For people prescribed IV infusions



Week 0 Week 2 Week 6

You will receive ENTYVIO by infusion at **Weeks 0, 2, and 6.**

For people prescribed self-administered injections



Week 0 Week 2

You will receive ENTYVIO by infusion at **Weeks 0 and 2.***

*Your healthcare provider will determine if it's appropriate to transition to self-administration with the ENTYVIO Pen after at least 2 IV doses. See [Page 4](#) for more information.

What should I do now?

Make sure to write down your appointment schedule and where you're getting your infusions.

Safety Considerations

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 7](#).



5

MAINTENANCE EVALUATION

After your initial IV starter doses, your healthcare provider may check in with you to see how the treatment is going.



If you plan to continue treatment with

IV INFUSIONS

[GO TO PAGE 5](#)



If you plan on transitioning to

SELF-ADMINISTERED INJECTIONS

[GO TO PAGE 6](#)

If you plan on transitioning to self-administered injection under the skin (subcutaneous), your healthcare provider will evaluate your situation and make a decision about the best time for you to transition.

Safety Considerations

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 7](#).




6

FIRST IV MAINTENANCE DOSE



8 weeks after your third IV infusion starter dose, you'll receive your first maintenance dose.



What should I do now?


Build an infusion day bag! A blanket, book, tablet, charger, snacks—whatever you need to feel comfortable during your infusions.

7

ONGOING IV MAINTENANCE



You will continue to receive ENTYVIO by IV infusion **every 8 weeks**.



- In the first year, you'll receive **8 doses of ENTYVIO**
- After the first year, you'll receive **6 doses each year**

What should I do now?

Stay in touch with your healthcare team about how you're feeling. Keeping a log of your symptoms in between visits can help you make the most of each conversation.

Safety Considerations

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 7](#).



6 RECEIVE SELF-INJECTION TRAINING KIT



This kit includes a training guide, practice pen, alcohol wipes, and more, all tucked inside a portable cooler you can use to take your medication on the go. Reach out to your *EntyvioConnect* support team to receive your training kit. If you haven't signed up for *EntyvioConnect* yet, see [Step 3](#) for more information.



Watch how to self-inject step by step

This video will give you a better sense of what it's like to self-inject ENTYVIO. Watch it at www.entyvio.com/how-to

7 FIRST SELF-INJECTION MAINTENANCE DOSE




After you receive training from your healthcare provider or *EntyvioConnect* Nurse Educator, you're ready to administer your first maintenance injection.

Make sure to read the full [Instructions for Use](#) included with your prescription.

What should I do now?

Set a reminder on your phone or calendar to inject ENTYVIO every two weeks.

8 ONGOING SELF-INJECTION MAINTENANCE



Every 2 Weeks

You'll continue to self-administer ENTYVIO **every 2 weeks, or up to 26 times per year.**

What should I do now?

Stay in touch with your healthcare team about how you're feeling. Keeping a log of your symptoms in between visits can help you make the most of each conversation.

Safety Considerations

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see **Important Safety Information** on [page 7](#).



Important Safety Information

- **Do not receive ENTYVIO® if you** have had an allergic reaction to ENTYVIO or any of its ingredients.
- **ENTYVIO may cause serious side effects, including:**
 - **Infusion-related and serious allergic reactions** can happen while you are receiving ENTYVIO or several hours after treatment. You may need treatment if you have an allergic reaction. Tell your healthcare provider or get immediate medical help if you get any of these symptoms during or after an infusion of ENTYVIO: rash, itching, swelling of your lips, tongue, throat or face, shortness of breath or trouble breathing, wheezing, dizziness, feeling hot, or palpitations (feel like your heart is racing).
 - **Infections.** ENTYVIO may increase your risk of getting a serious infection. Before receiving and during treatment with ENTYVIO, tell your healthcare provider if you think you have an infection or symptoms of an infection, such as fever, chills, muscle aches, cough, shortness of breath, runny nose, sore throat, red or painful skin or sores on your body, tiredness, or pain during urination.
 - **Progressive Multifocal Leukoencephalopathy (PML).** People with weakened immune systems can get progressive multifocal leukoencephalopathy (PML) (a rare, serious brain infection caused by a virus). Although unlikely while receiving ENTYVIO, a risk of PML cannot be ruled out. PML can result in death or severe disability. There is no known treatment, prevention, or cure for PML. Tell your healthcare provider right away if you have any of the following symptoms: confusion or problems thinking, loss of balance, change in the way you walk or talk, decreased strength or weakness on one side of the body, blurred vision, or loss of vision.
 - **Liver problems** can happen in people who receive ENTYVIO. Tell your healthcare provider right away if you have any of the following symptoms: tiredness, loss of appetite, pain on the right side of your abdomen, dark urine, or yellowing of the skin and eyes (jaundice).
- **The most common side effects of ENTYVIO include:** common cold, headache, joint pain, nausea, fever, infections of the nose and throat, tiredness, cough, bronchitis, flu, back pain, rash, itching, sinus infection, throat pain, pain in extremities, and with injections under the skin: pain, swelling, itching, hives, bruising, rash, or redness at the injection site. These are not all the possible side effects of ENTYVIO. Call your healthcare provider for medical advice about side effects.
- **Before receiving ENTYVIO, tell your healthcare provider about all of your medical conditions, including if you:** have or think you may have an infection or have infections that keep coming back; have liver problems; have tuberculosis (TB) or have been in close contact with someone with TB; have recently received or are scheduled to receive a vaccine; or if you are pregnant, breastfeeding, plan to become pregnant, or plan to breastfeed.
- **Tell your healthcare provider about all the medicines you take, especially if you take or have recently taken** Tysabri (natalizumab), Tyruko (natalizumab-sztn), a Tumor Necrosis Factor (TNF) blocker medicine, a medicine that weakens your immune system (immunosuppressant), or corticosteroid medicine.

Please see the full Prescribing Information, including the Medication Guide, for ENTYVIO and talk with your healthcare provider.

ENTYVIO is available for:

- **intravenous (IV) infusion: 300 mg vedolizumab.**
- **subcutaneous (SC) injection: 108 mg vedolizumab.**

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

What is ENTYVIO (vedolizumab)?

ENTYVIO is a prescription medicine used in adults for the treatment of:

- moderately to severely active ulcerative colitis (UC).
- moderately to severely active Crohn's disease (CD).

It is not known if ENTYVIO is safe and effective in children under 18 years of age.